



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed		for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details		
Display		Abstract	Show: 20	Sort	Send to		Text	

Entrez
PubMed

☐ 1: J Clin Microbiol 1997 Mar;35(3):570-7

Related Articles, Links

FREE full text article at
jcm.asm.org

A one-tube method of reverse transcription-PCR to efficiently amplify a 3-kilobase region from the RNA polymerase gene to the poly(A) tail of small round-structured viruses (Norwalk-like viruses).

PubMed
Services

Ando T, Monroe SS, Noel JS, Glass RI.

Viral Gastroenteritis Section, National Center for Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia 30333, USA.
txa5@CIDDDVD1.em.cdc.gov

Related
Resources

Amplification of a 3-kb genome region from the RNA polymerase gene to the 3' poly(A) tail of small round-structured virus (SRSV) by reverse transcription-PCR (RT-PCR) has been difficult to achieve because of a stable secondary structure in a region between the RNA polymerase gene and the 5' end of the second open reading frame. We have developed a one-tube RT-PCR method to efficiently amplify this region. The method comprises three procedures: purification of poly(A)+ RNA from a starting RNA solution by oligo(dT)30 covalently linked to latex particles, buffer exchange, and continuous RT and PCR in a single tube containing all reaction components. The key elements of this method are (i) first-strand cDNA synthesis with the Superscript II version of RNase H- Moloney murine leukemia virus reverse transcriptase at 50 degrees C for 10 min by using the RNA-oligo(dT)30 hybrid on the latex particles as the template and primer, and (ii) PCR by Taq and Pwo DNA polymerases mixed together with a mixture of 12 phased oligo(dT)25 antisense primers. The detection threshold of the one-tube RT-PCR method was as little as 0.2 ng of the crude RNA used as the source of the template. Using this method, we obtained 3-kb products from 24 SRSV strains previously characterized into four genetic groups. These included 5 P1-A, 4 P1-B, 5 P2-A, and 10 P2-B strains. Because SRSVs have not yet been cultivated in vitro, this novel method should facilitate molecular characterization of SRSVs to provide a firm scientific foundation for improvements and refinements of SRSV diagnostics.

PMID: 9041391 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Show:	20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	-------	----	---	------	---	---------	------	---

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Feb 19 2003 14:38:47

■



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed	▼ for longRT-PCR using Pwo and taq DNA polymerases						Go	Clear
Limits		Preview/Index		History		Clipboard		Details

One of your terms is not found in the database.

Entrez
PubMed

Display	Summary	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
Items 1-14 of 14								One page.	

PubMed
Services

☐ 1: [Held HA, Benner SA.](#)

[Related Articles](#), [Links](#)

Challenging artificial genetic systems: thymidine analogs with 5-position sulfur functionality.

Nucleic Acids Res. 2002 Sep 1;30(17):3857-69.

PMID: 12202771 [PubMed - indexed for MEDLINE]

☐ 2: [Attatippaholkun WH, Attatippaholkun MK, Nisalak A, Vaughn DW, Innis BL.](#)

[Related Articles](#), [Links](#)

A novel method for the preparation of large cDNA fragments from dengue-3 RNA genome by long RT-PCR amplification.

Southeast Asian J Trop Med Public Health. 2000;31 Suppl 1:126-33.

PMID: 11414441 [PubMed - indexed for MEDLINE]

☐ 3: [Mullan B, Kenny-Walsh E, Collins JK, Shanahan F, Fanning LJ.](#)

[Related Articles](#), [Links](#)

Inferred hepatitis C virus quasispecies diversity is influenced by choice of DNA polymerase in reverse transcriptase-polymerase chain reactions.

Anal Biochem. 2001 Feb 15;289(2):137-46.

PMID: 11161307 [PubMed - indexed for MEDLINE]

☐ 4: [Forrest R, Zhou H, Hickford JG.](#)

[Related Articles](#), [Links](#)

3'overhangs influence PCR-SSCP patterns.

Biotechniques. 2000 Nov;29(5):958-60, 962. No abstract available.

PMID: 11084854 [PubMed - indexed for MEDLINE]

☐ 5: [Kainz P.](#)

[Related Articles](#), [Links](#)

The PCR plateau phase - towards an understanding of its limitations.

Biochim Biophys Acta. 2000 Nov 15;1494(1-2):23-7.

PMID: 11072065 [PubMed - indexed for MEDLINE]

☐ 6: [Fromenty B, Demeilliers C, Mansouri A, Pessayre D.](#)

[Related Articles](#), [Links](#)

Escherichia coli exonuclease III enhances long PCR amplification of damaged DNA templates.

Nucleic Acids Res. 2000 Jun 1;28(11):E50.

PMID: 10871349 [PubMed - indexed for MEDLINE]

☐ 7: [Schulze E, Bettendorf M, Maser-Gluth C, Decker M, Schwabe U.](#)

[Related Articles](#), [Links](#)

Allele-dropout using PCR-based diagnosis for the splicing mutation in intron-2 of

Related
Resources

the CYP21B-gene: successful amplification with a Taq/Pwo-polymerase mixture.
Endocr Res. 1998 Aug-Nov;24(3-4):637-41.
PMID: 9888552 [PubMed - indexed for MEDLINE]

- ☐ 8: [Liu Q, Sommer SS.](#) Related Articles, Links

Subcycling-PCR for multiplex long-distance amplification of regions with high and low GC content: application to the inversion hotspot in the factor VIII gene.
Biotechniques. 1998 Dec;25(6):1022-8.
PMID: 9863056 [PubMed - indexed for MEDLINE]

- ☐ 9: [Iannelli F, Giunti L, Pozzi G.](#) Related Articles, Links

Direct sequencing of long polymerase chain reaction fragments.
Mol Biotechnol. 1998 Oct;10(2):183-5.
PMID: 9819816 [PubMed - indexed for MEDLINE]

- ☐ 10: [Abu Al-Soud W, Radstrom P.](#) Related Articles, Links

Capacity of nine thermostable DNA polymerases To mediate DNA amplification in the presence of PCR-inhibiting samples.
Appl Environ Microbiol. 1998 Oct;64(10):3748-53.
PMID: 9758794 [PubMed - indexed for MEDLINE]

- ☐ 11: [Gunther S, Sommer G, Von Breunig F, Iwanska A, Kalinina T, Sterneck M, Will H.](#) Related Articles, Links

Amplification of full-length hepatitis B virus genomes from samples from patients with low levels of viremia: frequency and functional consequences of PCR-introduced mutations.
J Clin Microbiol. 1998 Feb;36(2):531-8.
PMID: 9466771 [PubMed - indexed for MEDLINE]

- ☐ 12: [Rispeter K, Lu M, Lechner S, Zibert A, Roggendorf M.](#) Related Articles, Links

Cloning and characterization of a complete open reading frame of the hepatitis C virus genome in only two cDNA fragments.
J Gen Virol. 1997 Nov;78 (Pt 11):2751-9.
PMID: 9367360 [PubMed - indexed for MEDLINE]

- ☐ 13: [Ando T, Monroe SS, Noel JS, Glass RI.](#) Related Articles, Links

A one-tube method of reverse transcription-PCR to efficiently amplify a 3-kilobase region from the RNA polymerase gene to the poly(A) tail of small round-structured viruses (Norwalk-like viruses).
J Clin Microbiol. 1997 Mar;35(3):570-7.
PMID: 9041391 [PubMed - indexed for MEDLINE]

- ☐ 14: [Maruyama H, Kawakami H, Nakamura S.](#) Related Articles, Links

Reevaluation of the exact CAG repeat length in hereditary cerebellar ataxias using highly denaturing conditions and long PCR.
Hum Genet. 1996 May;97(5):591-5.
PMID: 8655136 [PubMed - indexed for MEDLINE]

Display	Summary	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
Items 1-14 of 14								One page.	

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

Feb 19 2003 14:38:47

■